

MOLD ARMOR®

DO IT YOURSELF MOLD TEST KIT TESTS

A minimum of 2 sample kits are needed to compare indoor to outdoor air quality. The first sample should be taken indoors where air quality is a concern and the second sample should be taken outdoors. If the amount of mold growing in the indoor sample is much greater than the outdoor sample, you may have a mold problem.

NOT INTENDED FOR PESTICIDAL USE

TO PREPARE THE PETRI DISH:

Remove the petri dish from the plastic bag and lay it on a level surface before removing lid. Remove the lid of the petri dish and pour the mold growth medium into the dish bottom. Replace the lid and swirl gently until the bottom is completely covered. Allow to gel for one hour on a level surface before removing lid. **Note:** You may only use one testing method per petri dish.

METHOD 1: INDOOR AIR QUALITY TEST

Place the prepared petri dish on a level surface in the location where you want to test for mold. Remove the lid of the petri dish and leave it open to the environment for 1 hour. After one hour replace the lid and allow to incubate for 48 hours. If there is no mold present within 48 hours, re-check the dish at 72 hours and 96 hours to be sure that no slow growing mold were overlooked. If mold growth is present after the incubation period, you may send the petri dish to our lab for analysis on the type of mold present. If no mold is present after 96 hours, no mold was detected.

METHOD 2: HVAC AIR SAMPLING (TAPE IS REQUIRED)

Mold can also come from your heating and air conditioning unit. To test, locate the air exhaust vent which is furthest away from the furnace. After preparing the petri dish, remove the lid and tape the open dish against the open air vent. Close all of the remaining air vents that are on the system. Turn the system fan on high for a 10 minute period, allowing forced air to hit the inside surface of the petri dish. After 10 minutes remove the petri dish, replace the lid, and allow to incubate for 48 hours. If no mold is present after 48 hours, re-check the dish at 72 hours and 96 hours to be sure that no slow growing mold were overlooked. If mold growth is present you may send the petri dish to our lab for analysis on the type of mold you have. If no mold is present after 96 hours, no mold was detected.

METHOD 3: VISUAL SAMPLING

Locate the visual mold growth. Take the sterile swab and wipe the tip across the surface mold growth. Be sure the mold adheres to the swab tip. Remove the lid from the petri dish. Very gently wipe the mold sample from the tip of the swab across the top surface of the medium on the petri dish, allowing the mold to adhere. Replace the lid. Be sure to send both the swab and the petri dish to our lab for analysis.

OPTIONAL LAB ANALYSIS (COST \$40)

See the information card found inside the envelope for instructions on the optional lab analysis.

Note: While Mold Armor Mold Test Kit can be an important part of protecting your family against mold, many factors contribute to any test's effectiveness. The Mold Test Kit can only detect airborne or surface mold spores (depending on test method) that are present in the area tested at the time of testing. If symptoms of mold exposure persist, a certified mold specialist should be consulted.

QUESTIONS? COMMENTS? CALL 888-328-7743

or visit www.moldarmor.com

WM Barr & Co., Inc., Memphis, TN 38113 / ©2010 WM Barr & Co., Inc.

Mold Armor® has stronger products for powerful mold stain control. Mold Armor Do-It-Yourself Mold Test Kit tests for the presence of mold within 48 hours. The kit can be used either indoors or outdoors. You will need one mold test kit for each area of concern. An optional Mail-In lab analysis (\$40) is available which identifies the type of mold present. (See directions for lab analysis on a prepaid lab mailer information card contained inside the package). The Mold Test Kit comes with a petri dish, a sterile swab and a bottle of mold growth medium.



WHY IS MOLD A CONCERN?
Mold produces mildew, musty odors, ugly stains and unpleasant odors. It grows on many surfaces and can heavily destroy your home both inside and outside. Over time, it can lead to an unstable living environment. Mold produces allergens which can cause allergic reactions and asthma attacks. It can also irritate the eyes, skin, nose, throat and lungs even among those with no allergies.

www.moldarmor.com

DO I HAVE A MOLD PROBLEM?

Use the Mold Armor® test kit to test for mold in your home or office. See our complete Mold Armor line of products to test, clean and prevent mold stain problems both indoors and outdoors.



MOLD ARMOR®

DO IT YOURSELF MOLD TEST KIT TESTS



MOLD ARMOR®



DO IT YOURSELF MOLD TEST KIT TESTS

- RELIABLE RESULTS IN AS LITTLE AS 48 HOURS
- SAFE AND EASY TO USE



INDOOR

OUTDOOR

OPTIONAL LAB ANALYSIS AVAILABLE

WHAT IS MOLD?

Mold and mildew are types of fungi. Mold is found both outdoors and indoors and can grow almost anywhere where there is water, high humidity, or damp conditions. Mold reproduces by means of tiny spores; the spores are invisible to the naked eye and float through outdoor and indoor air. There are approximately 10,000 mold spores on the head of a pin.

Mold may begin growing indoors when mold spores land on surfaces that are wet. Mold spores can be spread by vacuuming, walking on carpets or sitting on couches, for example. When checking for mold, look for signs of water damage, wet spots, staining, dampness or other evidence of water leaks or damage that may be a precursor to mold growth.